BoronMax

A Boron True Solution

Guaranteed Analysis:

Boron (B) 8.1%

Derived from boric acid

For use as a fertilizer in foliar and fertigation applications to supplement boron levels on agricultural crops.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application minimum of 5 gallons of water/acre; Dilute sprays minimum of 30 gallons of water/acre.
- Apply only with a minimum dilution of 1:100 in water.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- BoronMax is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Consult a NutriAg representative for application timing and necessity of its use.

Suggested Application Rates:

Сгор	Foliar Rates*	Fertigation Rates*
Forage Crops: Alfalfa, Clover, Hay Crops	Typical rates are	The number and
Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane	0.5 – 1 pint/acre/	frequency of
Legumes: Dry bean, Lentil, Peanut, Pea, Soybean	application	applications
Root Vegetables (including, but not limited to): Beet, Carrot,		determine
Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato,	Maximum rate	application rates.
Turnip	1 quart/acre/	
Leafy Vegetables (including but not limited to): Bok Choy,	application	Apply no more than
Celery,	with no more than	1 quart/acre/
Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach	2 quarts/season	application.
Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant,		
Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean,	Caution:	Apply no more than
Sweet Corn,	Crops vary widely	2 quarts/season
Tomato	in sensitivity to	
Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry,	boron.	Always apply toward
Gooseberry, Strawberry		the end of an
Tree Fruit (including but not limited to): Apple, Apricot,		irrigation cycle.
Avocado,		
Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum,		
Pomegranate, Prune, Quince		
Other Fruit: Citrus (all), Fig, Kiwi, Melons		
Nut Trees (including but not limited to): Almond, Filbert,		
Hazelnut, Pecan, Pistachio, Walnut		
Grapes: Currant, Raisin, Table, Wine		
Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower		



Danger

May damage fertility or the unborn child if swallowed. Contains 75-80% of an ingredient of unknown toxicity (Dermal and Inhalation)

Do not handle until all safety precautions have been read and understood. Wear protective gloves/protective clothing/eye protection/face protection.

If exposed or concerned: Get medical advice/attention.

Store locked up.

Dispose of contents/container in accordance with local, regional, national and international regulations

*Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or Net Contents: environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

CAUTION: The use of this fertilizing material on any crops other than those recommended may result in serious injury to

Caution: KEEP OUT OF REACH OF CHILDREN.

2.5 Gal. (9.46 L)

Net Weight:

26.48 lbs. (12.01 kg)

Net Weight/Gallon:

10.59 lbs. (4.80kg)

Batch No.: XXXXXXXXX