

A Copper True Solution with PAC* Chelation Technology

Guaranteed Analysis:

Copper (Cu) 4.2%

4.2% Chelated Copper

Derived from copper carbonate chelated with PolyAldoCarbosate.

For use as a fertilizer in foliar applications to supplement copper levels on agricultural crops.

*PAC refers to NutriAg's "PolyAldoCarbosate" Technology

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application and ground sprays minimum of 5 gallons of water/acre; Dilute sprays minimum of 30 gallons of water/acre.
- Apply only with a minimum dilution of 1:100 in water.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- CuMax is compatible with most agricultural chemicals and nutrients. Not compatible with aluminum equipment. Should be applied in conjunction with tissue and/or soil analysis.
- Consult a NutriAq representative for application timing and necessity of its use.

Suggested Application Rates:

Crop	Foliar Rates**
Forage Crops: Alfalfa, Clover, Hay Crops	
Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane	
Legumes: Dry bean, Lentil, Peanut, Pea, Soybean	
Root Vegetables (including, but not limited to): Beet,	
Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet,	
Sweet Potato, Turnip	
Leafy Vegetables (including but not limited to): Bok	0.5 – 4 pints/acre/
Choy, Celery,	application up to a
Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach	maximum of 4
Other Vegetables: Artichoke, Asparagus, Cucurbits,	applications/season
Eggplant, Okra, Onions (bunching, bulbing), Peppers,)
Rhubarb, Snap bean, Sweet Corn,	
Tomato	
Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry,	
Gooseberry, Strawberry	
Tree Fruit (including but not limited to): Apple, Apricot,	
Avocado,	
Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum,	
Pomegranate, Prune, Quince	
Other Fruit: Citrus (all), Fig, Kiwi, Melons	
Nut Trees (including but not limited to): Almond, Filbert,	
Hazelnut, Pecan, Pistachio, Walnut	
Grapes: Currant, Raisin, Table, Wine	
Other Crops: Herbs, Hops, Mint, Potato, Safflower,	
Spearmint, Stevia, Sunflower	

**Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

Caution: KEEP OUT OF REACH OF CHILDREN.



Danger

Fatal if swallowed. Causes severe skin burns and eye damage. Contains 28-31% of ingredients of unknown toxicity (Inhalation). Contains 17-19% of ingredients of unknown toxicity (Dermal).

Do not breathe dusts or mists. Wash hands and skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves/protective clothing/eye protection/face protection.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water/shower. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER/doctor. Specific treatment (see supplemental first aid instruction on this label). Wash contaminated clothing before reuse.

Store locked up.

Dispose of contents/container in accordance with local, regional, national and international regulations.

Net Contents:

2x2.5 Gal. (2x9.46 L)

Net Weight:

46.54lbs. (21.12 kg)

Net Weight/Gallon: 9.31 lbs. (4.22 kg)

Batch No.: XXXXXX