Enviro Zinc

OMRI LISTED For Organic Use





A zinc true solution for use in organic production.

Guaranteed Analysis:

Zinc (Zn) 8.5% Derived from zinc sulfate

For use in organic production as a fertilizer in foliar, fertigation and soil applications to supplement zinc levels on agricultural crops. Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Directions for Use (Use only as Directed):

- · Avoid foliar application in extreme weather conditions such as high temperatures and poor drying conditions.
- Aerial application minimum of 5 gallons of water/acre; Dilute sprays minimum of 20 gallons of water/acre or apply with sufficient water to deliver spray uniformly on leaf surfaces.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- Enviro Zinc is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- Consult a NutriAg representative for application timing and necessity of its use.

Suggested Application Rates:

CROP	FOLIAR RATES**	FERTIGATION RATES**	SOIL RATES**
Forage Crops: Alfalfa, Clover, Hay Crops	1-3 pints/acre/	The number	This product
Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane	application	and frequency	is compatible
Legumes: Dry bean, Lentil, Peanut, Pea, Soybean		of applications	with many liquid
Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip,		determine	fertilizers.
Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip		application rates.	
Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro,			Fertilizers vary
Cole Crops, Lettuce, Parsley, Spinach		Apply no more than	in analysis and
Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions		2 quarts/acre/	quality, and
(bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato		application.	should be jar
Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry			tested with this
Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry,		Apply no more than	product before
Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince		8 quarts/season.	use.
Other Fruit: Citrus (all), Fig, Kiwi, Melons			
Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan,		Always apply	Apply up to
Pistachio, Walnut		toward the end of an	2 quarts/acre at
Grapes: Currant, Raisin, Table, Wine		irrigation cycle.	planting or as a
Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower			side dress.

**Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg Ltd. warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg Ltd. NutriAg Ltd. is not liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

Rev: 080317CA

Net Contents: Net Weight: Net Weight/Gallon: Batch No.:

DANGER

Causes serious eye damage. Contains 24.5% of ingredient(s) of unknown toxicity (Dermal and Inhalation)

Wear eye protection/face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE/doctor.

Dispose of contents/ container in accordance with local, regional, national and international regulations.

This product contains zinc (Zn) and may be used for fertility only with a documented zinc. The use of the fertilizing material on any crops other than those recommended may result is serious injury to crop(s).

KEEP OUT OF REACH OF CHILDREN.

NutriAg Ltd.

39 Gail Grove, Toronto, ON, Canada, M9M 1M5. Tel: (416) 636-1555

