

ZinManMax

A Zinc, Manganese, Molybdenum and Boron True Solution with PAC* Chelation Technology

Guaranteed Analysis:

Boron (B)	0.5%
Manganese (Mn)	2.7%
Molybdenum (Mo)	0.25%
Zinc (Zn)	5.1%

Derived from boric acid, manganese sulphate, sodium molybdate and zinc oxide with PolyAldoCarbosate

For use as a fertilizer in foliar, fertigation and soil applications to supplement micronutrient levels on agricultural crops.

***PAC refers to NutriAg's "PolyAldoCarbosate" Technology**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application and ground sprays – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- ZinManMax is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- **Consult a NutriAg representative for application timing and necessity of its use.**

Suggested Application Rates:

Crop	Foliar Rates**	Fertigation Rates**	Soil Rates**
Forage Crops: Alfalfa, Clover, Hay Crops Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane Legumes: Dry bean, Lentil, Peanut, Pea, Soybean Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince Other Fruit: Citrus (all), Fig, Kiwi, Melons Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut Grapes: Currant, Raisin, Table, Wine Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower	0.5 – 3 quarts/acre/ application	The number and frequency of applications determine application rates. Apply no more than 3 quarts/acre/ application. Apply no more than 9 quarts/season Always apply toward the end of an irrigation cycle.	This product is compatible with many liquid fertilizers. Fertilizers vary in analysis and quality, and should be jar tested with this products before use. Apply up to 3 quarts/acre at planting or as a side dress



Danger

Causes serious eye damage. Causes skin irritation. Suspected of causing genetic defects. May cause damage to organs through prolonged or repeated exposure. Contains 18-22% of ingredients of unknown toxicity (Inhalation and Dermal). Contains 2-3% of ingredients of unknown toxicity (Oral).

Do not breathe mist, vapours, spray. Wash hands and skin thoroughly after handling. Wear protective gloves. Wear eye protection/face protection.

IF ON SKIN: Wash with plenty of water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER/doctor. Get medical advice/attention if you feel unwell. Specific treatment (see supplemental first aid instruction on this label). If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.

Dispose of contents/container in accordance with local, regional, national and international regulations.

Caution: KEEP OUT OF REACH OF CHILDREN.

**Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg Ltd. warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg Ltd. NutriAg Ltd. will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

CAUTION: The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crops.

CAUTION: The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

Caution: KEEP OUT OF REACH OF CHILDREN.

Net Contents:
2x2.5 Gal. (2x9.46 L)

Net Weight:
51.46 lbs. (23.34 kg)

Net Weight/Gallon:
10.29 lbs. (4.67 kg)

Batch No.: XXXXX