

# TrioMax

## A Zinc, Manganese, Copper and Molybdenum True Solution with PAC\* Chelation Technology

### Guaranteed Analysis:

Copper (Cu)	1.4%
Manganese (Mn)	2.9%
Molybdenum (Mo)	0.24%
Zinc (Zn)	1.4%
Boron (B)	0.28%

For use as a fertilizer in foliar, fertigation and soil applications to supplement zinc, manganese, copper, molybdenum and boron levels on agricultural crops.

\*PAC refers to NutriAg's "Polyaldocarbosate" Chelation Technology

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

### Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- TrioMax is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis. Not compatible with aluminum equipment.
- **Consult a NutriAg representative for application timing and necessity of its use.**

### Suggested Application Rates:

Crop	Foliar Rates**	Fertigation Rates**	Soil Rates**
<b>Forage Crops:</b> Alfalfa, Clover, Hay Crops <b>Field Crops:</b> Cereals, Corn, Cotton, Sorghum, Sugar cane <b>Legumes:</b> Dry bean, Lentil, Peanut, Pea, Soybean <b>Root Vegetables (including, but not limited to):</b> Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip <b>Leafy Vegetables (including but not limited to):</b> Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach <b>Other Vegetables:</b> Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato <b>Berry Crops:</b> Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry <b>Tree Fruit (including but not limited to):</b> Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince <b>Other Fruit:</b> Citrus (all), Fig, Kiwi, Melons <b>Nut Trees (including but not limited to):</b> Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut <b>Grapes:</b> Currant, Raisin, Table, Wine <b>Other Crops:</b> Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower	1-4 pints/acre/ application	The number and frequency of applications determine application rates.  Apply no more than 2 quarts/acre/ application.  Apply no more than 8 quarts/season.  Always apply toward the end of an irrigation cycle.	This product is compatible with many liquid fertilizers.  Fertilizers vary in analysis and quality, and should be jar tested with this product before use.  Apply up to 2 quarts/acre as a side dress.



#### Danger

Fatal if swallowed. Contains 19-23% of ingredients with unknown toxicity (Inhalation). Contains 16-19% of ingredients with unknown toxicity (Dermal). Contains 12-14% of ingredients with unknown toxicity (Oral)

Wash hands and skin thoroughly after handling. Do not eat, drink or smoke when using this product.

IF SWALLOWED: Immediately call a POISON CENTER/doctor. Specific treatment (see supplemental first aid instruction on this label). Rinse mouth.

Store locked up.

Dispose of contents/container in accordance with local, regional, national and international regulations.

CAUTION: This product contains Boron (B). The use of this fertilizing material on any crops other than those recommended may result in serious injury to the crops.

CAUTION: The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

\*\*Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.

**Warranty Statement:** Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg Ltd. warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg Ltd. NutriAg Ltd. will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

**CAUTION:** This fertilizing material contains boron. The use of the fertilizing material on any crops other than those recommended may result in serious injury to crop(s). **Caution: KEEP OUT OF REACH OF CHILDREN.**

**Net Contents:**  
2.5 Gal. (9.46 L)

**Net Weight:**  
27.1 lbs. (12.3 kg)

**Net Weight/Gallon:**  
10.8 lbs. (4.92 kg)

**Batch No.:** XXXXXX