

TruPhos ManZin 6-20-5

Guaranteed Analysis:		
Total Nitrogen (N)	6%	Sulphur (S) 1.8%
Available Phosphoric Acid (P ₂ O ₅)	20%	Manganese (Mn) 3%
Soluble Potash (K ₂ O)	5%	3% Water Soluble Manganese
		Zinc (Zn) 1%

An OrthoPhosphate True Solution with Manganese and Zinc

Derived from orthophosphate, potassium carbonate, urea, sulfates of manganese, zinc oxide.

For use as a fertilizer in foliar and fertigation applications to supplement phosphorus, potassium, nitrogen, sulfur, manganese and zinc levels on agricultural crops.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Directions for Use (Use only as Directed):

- Avoid foliar application in poor drying conditions such as very high humidity, heavy dew set or continually rainy weather.
- Do not foliar apply when ambient temperatures are expected to exceed 85°F before application completely dries on foliage.
- Aerial application and ground sprays – minimum of 5 gallons of water/acre; Dilute sprays – minimum of 30 gallons of water/acre.
- Exercise caution when applying to any crop during bloom, as many factors can affect fruit set and final fruit quality.
- TruPhos ManZin is compatible with most agricultural chemicals and nutrients. Should be applied in conjunction with tissue and/or soil analysis.
- **Consult a NutriAg representative for application timing and necessity of its use.**

Suggested Application Rates:

Crop	Foliar Rates*	Fertigation Rates*
Forage Crops: Alfalfa, Clover, Hay Crops Field Crops: Cereals, Corn, Cotton, Sorghum, Sugar cane Legumes: Dry bean, Lentil, Peanut, Pea, Soybean Root Vegetables (including, but not limited to): Beet, Carrot, Garlic, Parsnip, Radish, Rutabaga, Sugar Beet, Sweet Potato, Turnip Leafy Vegetables (including but not limited to): Bok Choy, Celery, Chard, Cilantro, Cole Crops, Lettuce, Parsley, Spinach Other Vegetables: Artichoke, Asparagus, Cucurbits, Eggplant, Okra, Onions (bunching, bulbing), Peppers, Rhubarb, Snap bean, Sweet Corn, Tomato Berry Crops: Blueberry, Caneberry, Cranberry, Elderberry, Gooseberry, Strawberry Tree Fruit (including but not limited to): Apple, Apricot, Avocado, Cherry, Nectarine, Olive, Peach, Pear, Persimmon, Plum, Pomegranate, Prune, Quince Other Fruit: Citrus (all), Fig, Kiwi, Melons Nut Trees (including but not limited to): Almond, Filbert, Hazelnut, Pecan, Pistachio, Walnut Grapes: Currant, Raisin, Table, Wine Other Crops: Herbs, Hops, Mint, Potato, Safflower, Spearmint, Stevia, Sunflower	1 quart/acre/ application Apply no more than 3 quarts/season	The number and frequency of applications determine application rates. Apply no more than 1 quart/acre/application. Apply no more than 3 quarts/season. Always apply toward the end of an irrigation cycle.



Warning

Causes skin irritation. Causes serious eye irritation. Suspected of causing genetic defects. May cause cancer. Contains 47-51% of ingredients with unknown toxicity (Inhalation). Contains 21-33% of ingredients with unknown toxicity (Dermal). Contains 9-11% of ingredients with unknown toxicity (Oral).

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wash hands and skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.

IF ON SKIN: Wash with plenty of water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Specific treatment (see supplemental first aid instructions on this label). If skin irritation occurs: Get medical advice/attention. If eye irritation persists: Get medical advice/attention. Take off contaminated clothing and wash it before reuse.

Store locked up.

Dispose of contents/container in accordance with local, regional, national and international regulations.

***Application rates vary according to specific crop requirements, management objectives, soil nutrient availability and/or environmental conditions. Lower rates are recommended for general crop maintenance or when applying multiple times per season; higher rates are recommended when applying a minimum number of applications to crops that require repeat applications, or where more severe deficiency is present. Higher volumes of water should be used for higher rates of fertilizer. For application timing charts, please consult a product manual. Consult your NutriAg representative for specific crop recommendations.**

Warranty Statement: Sellers guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on this condition. NutriAg warrants that the product conforms to the chemical description on the label and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product such as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of NutriAg. NutriAg will not be liable for consequential, special or indirect damages resulting from the use of or handling of this product. All such risks shall be assumed by the buyer.

Caution: KEEP OUT OF REACH OF CHILDREN.

Net Contents:
2.5 Gal. (9.46 L)

Net Weight:
29.49 lbs. (13.38 kg)

Net Weight/Gallon:
11.80 lbs. (5.35 kg)

Batch No.: XXXXXX